

CDC Guidance - Vaccine Coadministration

COVID-19 vaccines were previously recommended to be administered alone, with a minimum interval of 14 days before or after administration of any other vaccines. This was out of an abundance of caution and not due to any known safety or immunogenicity concerns. However, substantial data have now been collected regarding the safety of COVID-19 vaccine currently authorized by FDA for use under EUA. Although data are not available for COVID-19 vaccines administered simultaneously with other vaccines, extensive experience with non-COVID-19 vaccines has demonstrated that immunogenicity and adverse event profiles are generally similar when vaccines are administered simultaneously as when they are administered alone.

COVID-19 vaccines and other vaccines **may now be administered without regard to timing**. This includes simultaneous administration of COVID-19 vaccine and other vaccines on the same day, as well as coadministration within 14 days. It is unknown whether reactogenicity of COVID-19 vaccine is increased with coadministration, including with other vaccines known to be more reactogenic, such as adjuvanted vaccines or live vaccines. When deciding whether to coadminister another vaccine(s) with COVID-19 vaccine, providers should consider whether the patient is behind or at risk of becoming behind on recommended vaccines, their risk of vaccine-preventable disease (i.e., during an outbreak or occupational exposures), and the reactogenicity profile of the vaccines.

If multiple vaccines are administered at a single visit, administer each injection in a different injection site. For adolescents and adults, the deltoid muscle can be used for more than one intramuscular injection.

Best practices for multiple injections include:

- Label each syringe with the name and the dosage (amount) of the vaccine, lot number, the initials of the preparer, and the exact beyond-use time, if applicable.
- Separate injection sites by 1 inch or more, if possible.
- **Administer the COVID-19 vaccines and vaccines that may be more likely to cause a local reaction (i.e., tetanus-toxoid-containing and adjuvanted vaccines) in different limbs, if possible.**

<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html#Coadministration>

May 14, 2021

This work is licensed under the Creative Commons Attribution-NoDerivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nd/4.0/>

This policy and procedure is not intended to replace the informed judgment of individual physicians, nurses or other clinicians nor is it intended as a statement of prevailing community standards or minimum standards of practice. It is a suggested method and technique for achieving optimal health care, not a minimum standard below which residents necessarily would be placed at risk.